



REVIEW ARTICLE

Positive pressure ventilation: what is the real cost?

N. Soni* and P. Williams

Imperial College Medical School, Chelsea and Westminster Hospital, 369 Fulham Road,
London SW10 9NH, UK

*Corresponding author. E-mail: n.soni@imperial.ac.uk

Positive pressure ventilation is a radical departure from the physiology of breathing spontaneously. The immediate physiological consequences of positive pressure ventilation such as haemodynamic changes are recognized, studied, and understood. There are other significant physiological interactions which are less obvious, more insidious, and may only produce complications if ventilation is prolonged. The interaction of positive pressure with airway resistance and alveolar compliance affects distribution of gas flow within the lung. The result is a wide range of ventilation efficacy throughout different areas of the lung, but the pressure differentials between alveolus and interstitium also influence capillary perfusion. The hydrostatic forces across the capillaries associated with the effects of raised venous pressures compound these changes resulting in interstitial fluid sequestration. This is increased by impaired lymphatic drainage which is secondary to raised intrathoracic pressure but also influenced by raised central venous pressure. Ventilation and PEEP promulgate further physiological derangement. In theory, avoiding these physiological disturbances in a rested lung may be better for the lung and other organs. An alternative to positive pressure ventilation might be to investigate oxygen supplementation of a physiologically neutral and rested lung. Abandoning heroic ventilation would be a massive departure from current practice but might be a more rationale approach to future practice.

Br J Anaesth 2008; **101**: 446–57

Keywords: complications, respiratory; lung, compliance; lung, pathophysiology; ventilation, positive end-expiratory pressure

Positive pressure ventilation during the Copenhagen polio epidemic was the key to the reduction in mortality from 87% to 40%. It was achieved using medical students not sophisticated ventilators, with minimal understanding of the physiology of gas exchange, and with no real means of monitoring either oxygenation or carbon dioxide. It rapidly became obvious that its use could be extrapolated to the management of respiratory failure.^{72 73 144} In the first edition of *Ventilation of the Lungs*, Mushin comments that ‘positive pressure ventilation is a considerable deviation from the normal physiological mechanism of respiration’. Implicit in this statement is the notion that there will be a pathophysiological price to pay for disturbing the normal physiology of breathing. The immediate interactions, such as haemodynamic changes, are well delineated, but it is increasingly clear that not all the complications are either obvious or immediate. The ARDSnet study exposed previously concealed mortality suggesting the insidious and late impact of shear forces and other inflammatory reactions secondary to mechanical ventilation.¹

The driving force of ventilatory techniques in the intensive care unit (ICU) is to achieve oxygenation which in severe lung injury may be both difficult and prolonged.

This review revisits the physiological cost involved in achieving oxygenation using positive pressure ventilation in the critically ill, and discusses some of the possible interactions (Table 1).

Oxygenation and ventilation

Each ICU has rules as to what constitutes acceptable oxygenation and this varies with the patient and circumstances.¹⁶ There are many examples of people functioning reasonably with far lower oxygenation indices than would be acceptable in an ICU. These range from high altitude residents and mountaineers to patients living with chronic airways disease.⁹⁸ In the ICU, monitoring of oxygenation still focuses on inspired oxygen concentration and arterial blood gases to indicate adequacy, whereas in other areas of practice, emphasis has moved to surrogates of tissue hypoxia such as ScVO₂ and lactate (Fig. 1). With

Table 1 Some areas of potential physiological interaction with positive pressure ventilation

Positive pressure effects on intrathoracic pressures
Pressure and distribution of airflow into the alveoli
Pressure, stretch, and the lung
Surfactant functions
Positive pressure and cardiac output
Pulmonary capillary blood flow
Positive pressure and the lymphatics
Positive pressure and organ system function




Fig 1 Monitoring oxygenation. Based on arterial values which correlate poorly with adequacy of tissue oxygenation.

circulation, tissue and cell transport, and mitochondrial function interposed between lung and cell, the arterial values of oxygenation may not be the ideal monitoring parameters.

In the critically ill, several observations regarding oxygenation can be made.

- (i) Even with relatively low arterial saturations, there may be adequate tissue oxygenation.
- (ii) When tissue oxygenation is in jeopardy, it is very often due to problems not related directly to the lungs.
- (iii) Saturation and Pa_{O_2} below the acceptable normal range correlate poorly with tissue oxygenation.
- (iv) When patients are difficult to oxygenate, tolerance of lower values develops rapidly, provided there are no markers of tissue hypoxia.
- (v) It is often stated that ‘few patients with lung injury die of hypoxaemia’. Clearly, some do but often death is due to acquired complications.

Margins of safety for oxygenation are essential, but difficulty arises when it is necessary to use aggressive ventilation to achieve margins that may be far in excess of actual requirement. If this is associated with late complications, then methods of assessing adequacy of oxygenation may need to be reconsidered.

Breathing, ventilation, and intrathoracic pressures

In inspiration, small negative intrapleural, interstitial, and alveolar pressures are generated allowing the lung to inflate. On expiration, intrapleural pressure returns towards

atmospheric, but remains negative, whereas interstitial and alveolar pressures return towards atmospheric or slightly positive pressure.

Positive pressure ventilation reverses this cycle with high intrathoracic pressures on inspiration which are transmitted, to the alveoli and the interstitial tissues, following the path of least resistance to inflate the alveoli. The intrapulmonary and interstitial pressures remain positive throughout inspiration but will return towards atmospheric pressure on expiration, unless PEEP is added when the pressures remain positive throughout.

Ventilation, alveolar ventilation, and recruitment

During spontaneous breathing, the negative pressure gradient generates air distribution which is locally influenced by the state of the airways, alveoli, and interstitium. Surfactant modifies the effects of Laplace’s law at these low pressures so that small alveoli are both relatively easy to inflate and have less tendency to collapse on expiration. Despite these effects, there is regional variation in gas volume distribution even in volunteers breathing normally.⁸⁹ The same factors influence flow with positive pressure ventilation, but the forces acting are different. In the tracheal tube, a single pressure applied over a specific time produces a waveform which forces gas through the airways. The state of the airways, in particular their resistance, and the alveolar compliance determine the effect of that pressure in different regions of the lung. These factors combined can be considered as the individual ‘time constants’ of the lung regions or of individual alveoli. In damaged lung, the range of different time constants between regions will increase even if the surfactant was normal. Positive pressure will preferentially aerate high compliance areas at the expense of lower compliance areas, whereas collapsed alveoli may require high sustained pressure to force them open, but this may be ineffective. It is a daily observation in the operating theatre that re-inflation of a collapsed but previously normal lung requires sustained high positive pressure and it is a gradual and far from uniform process (Fig. 2).

Recruitment is the opening and maintaining open potentially under ventilated areas. The fundamental concept behind recruitment is to use the physics of inflation judiciously, to increase the number of alveoli and hence alveolar surface area involved in gas exchange. The central tenet is that this will improve oxygenation. A wide range of strategies aimed to ‘recruit’ alveoli have been advocated, but the problem with any of these techniques is that the variation in time constants between alveoli and lung regions precludes a single pressure or waveform being suitable for all lung regions.

CT scanning and electrical impedance tomography (EIT) have revolutionized the ability to ‘see’ what actually happens to lung inflation with positive pressure ventilation.

Using EIT, the inferred pressure–volume curve clearly shows a lower and upper inflection point (increased/reduced conductivity). The lower inflection point is thought to represent recruitment whereas the upper inflection point represents over-distension, although this is contentious (Fig. 3). Physiologically, this methodology has clearly shown not only regional variation in both recruitment and over-distension but that pathology in the lung increases the diversity of these changes.³⁴ EIT indicates a huge regional variation in both recruitment and over-inflation, in injured lungs, both within and between lungs.^{18 58–60 133} It also demonstrates that whole lung inflection points do not represent individual areas.

CT has also been used to assess recruitment. In ARDS, a recruitment technique using an initial sustained high pressure with subsequent PEEP at various levels to maintain recruitment only achieved an estimated mean recruitment of 13%, and showed that a large part of the lung was not recruitable.^{48 65} It did indicate that high levels of PEEP, 15 mm Hg, are more effective than lower levels, 5 mm Hg, both in maintaining alveolar patency and improving oxygenation. PEEP is also effective in preventing

collapse and derecruitment.^{43 87 130} All these techniques result in sustained increase in intrathoracic pressure.

A further consideration is that pressure is effectively distributed across compliant lung but dampened by poorly compliant areas. Paradoxically, the techniques used to open closed airways, probably results in increased gas distribution not to the abnormal areas but rather to the functional areas over-inflating them and potentially impairing their function.^{17 127} The result of recruitment is therefore unpredictable.

In summary, the physics of the lung inflated by a single pressure waveform will result in unpredictable gas distribution due to variable regional compliance. There will be over-distension of normal alveoli, and although there will be some recruitment, it may be very limited, especially in areas of collapse or consolidation. The combination of high inspiratory pressures and high levels of PEEP will result in sustained pressures throughout the respiratory cycle.

Recruitment in the short term with relatively normal lungs works well but in damaged lungs, and particularly over time, recruitment techniques may exacerbate problems while apparently improving oxygenation indices.




Fig 2 Alveolar inflation. (A) Represents the beginning of inspiration. Pressure P acts on a consolidated alveolus, a normal alveolus and a distended alveolus. (B) End inspiration. Pressure P produces a broad spectrum of effects.

Ventilation pressure and stretch

The reduction in mortality highlighted by the ARDSnet study was associated with lower peak pressure ventilation, but it also appears that the shear forces that occur particularly in initiating inflation may cause injury.^{1 20 24 36 49 85 121 122 124 149}

Stress and shear forces exerted during the cyclic opening and closing of alveoli result in increased cytokine production and white cell sequestration, as does over-distension of alveoli.^{19 20} These effects are seen over a range of ventilatory patterns and it may predispose to injury and infection.^{78 128 143} Repetitive identical ventilation exerts different




Fig 3 Pressure–volume curve and the effects on alveoli.

and more damaging shear forces than variable ventilation where both volume and duration are varied.^{13 95} These inflammatory effects originating in the lung may cause secondary inflammatory problems in other organs.^{35 38 68 84 92}

In recruitment, manoeuvres the lowering of the peak pressure is important whereas the use of higher PEEP maintains alveolar patency and reduces the shear forces. The net effect is high pressure throughout the ventilatory cycle.

Ventilation and surfactant

Surfactants modify the surface tension and hence the law of Laplace. The denser the molecules, the lower the surface tension which enables small alveoli, with dense surfactant, to remain open but also facilitates inflation during inspiration. In larger alveoli during inflation, the density of surfactant decreases as the surface area increases and the influence on the surface tension decreases, so that the surface tension increases. This is highly efficient in spontaneous breathing at low pressures as it facilitates effective gas distribution throughout the lung. With positive pressure ventilation, air is forced into the lung under pressure and the resultant pressure waveform may generate wave formation in the surfactant layers altering the uniformity of spread. This is known in surfactant technology, but there is little or no information relating to surfactant wave behaviour in the lung. A fundamental difference in the performance of surfactant between spontaneous and positive pressure ventilation may be due to distortion of surfactant spread. This is unlikely to be uniform across the lung due to wide variation in pressure and therefore of flow.

Positive pressure also influences the production and function of surfactant. In lung injury, type II pneumocytes are reduced in number and surfactant production is reduced.⁹⁹ Function is altered by the mix of cholesterol and protein which then affects the ability of the surfactant to form surface structures.^{83 99 137} This is exacerbated by inflammation which releases protein and other materials that mix with the surfactant, mediators that directly affect surfactant, membrane permeability changes that allow fluid dilution of surfactant, and polymerizing fibrin that adsorbs these surface active compounds, effectively removing them.^{115 116 126 129 131 138} Infection is implicated, with evidence that both pseudomonas and pneumocystis pneumonias alter surfactant function, although it is uncertain whether it is due to the disease or the ventilation. It is likely other infections do likewise.^{40 113 136} Medical intervention such as lavage effectively removes surfactant impairing regional and whole lung compliance.²⁹ Surfactant dysfunction changes compliance and exaggerates heterogeneous lung inflation necessitating ventilation patterns which may cause further damage.^{11 30 69 123}

Surfactant has a role in the immune defences of the lung. The lung provides more than 50 m² of contact across a fine membrane and is therefore a potential site for invasion. Not

only does surfactant influence inflammatory response, but it has a substantial role in mucosal immunity and at least two surfactants have activity as collectins (molecules that bind to foreign particles such as bacteria, fungi, and viral particles and assist in their phagocytosis).^{21 50 53 54 67}

In conclusion, both positive pressure and lung injury alter the production and function of surfactant. Changes in surface tension and loss of immunocompetence both predispose to lung injury and infection. A vicious cycle ensues where the disease and the supportive treatment may be synergistic in producing deleterious effects and will make positive pressure ventilation both more difficult and more prone to cause further injury.

Effect on the cardiovascular system

The effects of positive pressure ventilation on the cardiovascular system are well known. In normal breathing, the negative pressure phase of inspiration assists venous return, alleviates pressure on the pulmonary capillaries, and encourages flow. With positive pressure ventilation, the intrathoracic pressure increases during inspiration causing a decrease in venous return, right ventricular output, and pulmonary blood flow. Paradoxically, there may be a reduction in right ventricular impedance, but whether this offsets the decrease in venous return is unknown.⁹⁰ There is some evidence to show that as PEEP increases, right ventricular outflow tends to decrease, reducing return to the left ventricle.^{91 134} This effect may be partially offset by the increased intrathoracic–extrathoracic gradient assisting left ventricular output. This may be beneficial in cardiac failure but is not usually relevant in other circumstances where raised intrathoracic pressure reduces output.¹⁰²

On expiration, the intrathoracic pressure returns towards zero so that venous return will increase. If PEEP is applied, positive intrathoracic pressure continues to inhibit venous return through expiration. The reduced collapsibility of the inferior vena cava often seen with ventilation and PEEP is a clear indicator of a degree of venous stasis.⁹¹ Fluid administration improves venous return and cardiac output but at the cost of increased central venous pressures (CVP) and hence increased end-capillary pressures in the lungs and other organs.

Positive pressure ventilation is associated with salt and water retention. While classically this is due to increased secretion of anti-diuretic hormone mediated through the reduced atrial distension seen with positive pressure ventilation, more recently atrial natriuretic peptide has been implicated.^{4 42 66} Correction of this by intravascular fluid administration results in further fluid retention.^{5 7 139}

The pulmonary capillary and blood flow

The mean capillary pressure is usually 7–10 mm Hg, a value derived from the gradient between the pulmonary

artery pressure and the venous outflow pressure in the pulmonary vein. Intrathoracic pressure will act uniformly across the capillary and should not directly influence gradient, unless the capillary is physically compressed by the raised interstitial pressure along its course.

During normal breathing, the pressures in the interstitium and the alveoli are small relative to the capillary perfusion pressure. This low pressure system enables the hydrostatic and oncotic pressures to facilitate fluid and substrate movement between compartments but also ensures unimpeded blood flow. Furthermore, the pressures within the lung generally are lower than capillary pressure in all phases and do not influence capillary flow. The exceptions being areas higher than the atrium where the pressure gradients are lower, and in the dependent areas where, theoretically, lung tissue weight may press on capillaries. These effects are small and transient. In disease, this can be different. Patients with chronic obstructive pulmonary disease with hyperinflation have a high intrathoracic pressure on expiration which compresses the interstitium and the capillaries, thus increasing capillary resistance and modifying blood flow.⁸⁸ On inspiration, as the intrathoracic pressures decrease, flow can recover.

Positive pressure ventilation in a critically ill patient, even with peak inspiratory pressures limited to 30 mm Hg, produces interstitial pressures considerably in excess of capillary pressure and will tend to compress the capillary and impede flow. In expiration, PEEP of 15 mm Hg may still exceed capillary pressure and prevents recovery of normal flow.⁹⁷ Compensatory mechanisms will also be modified. Hypoxic vasoconstriction, which is pre-capillary and which is highly effective in diverting blood flow during spontaneous breathing at low pressures, may be offset by the magnitude of the external pressures being applied.¹⁴⁰ The capillary pressure itself increases minimally with pre-capillary vasoconstriction, probably in the order of 1 mm Hg; hence, the capillary will remain susceptible to external pressure.^{23 120} At high altitude, some individuals respond to hypoxia by a more generalized, albeit still regionally heterogeneous, increase in pulmonary artery pressure which influences flow, presumably diverting it to appropriate areas.^{32 64} During positive pressure ventilation, compliant inflatable lung regions readily transmit pressure and in these areas the pressure will impede capillary flow whereas non-compliant damaged or infected lung will dampen pressure transmission and may have better perfusion. Compensatory diversion is unlikely to overwhelm the effects of this magnitude. Perfusion-ventilation mismatch is inevitable.

The high venous pressures, secondary to positive pressure ventilation, will also adversely affect the pressure gradients affecting Starling's forces and encouraging fluid flux, leading to interstitial fluid retention.

In the critically ill, increasing pulmonary artery pressures, high inflation pressures, and persistently high venous pressures probably damage the integrity of the capillary endothelium. 'Capillary stress failure' or disruption of the

capillary is seen in extreme exercise in racehorses and is probably also seen in humans with high pulmonary artery pressures, high alveolar pressure, and in altitude hypoxia.^{64 132 143}

Capillary perfusion pressure is of central importance to lung function. The consequences of persistently raised intrathoracic pressure are altered flow by compression of the capillary, modified Starling's forces, and raised end-capillary pressures. Compensatory mechanisms are unlikely to be effective and may aggravate ventilation-perfusion mismatch.

Lymphatics

The lymphatics subserve at least two functions, drainage and defence. The lung has a massive surface area continuously exposed to a large volume of poorly filtered air. Fluid is drained from the interstitium by the reticuloendothelial system that removes extraneous material before it reaches the circulation.

The lung lymphatics consist of thin, single cell, conduits within the interstitium with valves to assist unidirectional flow. During inspiration, the negative pressure in the lung is transmitted to the interstitium and to the lymphatics and a gradient allows fluid to drain into the lymphatics. This is highly efficient and lymph flow can increase considerably if needed.^{9 41 51 52 76} As the lymphatics fill, pressure increases and elasticity increases forward flow aided by muscle activity in the limbs or by ventilatory movement.² Towards the hilum, smaller channels coalesce into larger ones and they end in the thoracic duct or large lymphatics which drain into central veins.¹¹²

The pressure in the peripheral lymphatics reaches a maximum around 4 mm Hg, at which flow is maximal but if the pressure continues to increase the walls overstretch and leak. This pressure of around 4 mm Hg is adequate to provide a hydrostatic gradient between the lymphatics and the central veins which have very low pressures during the inspiration. The collecting ducts are slightly more robust and contain smooth muscle. This helps the lymphatics compensate for some outflow resistance, but they are still easily compressed. Flow will be relatively easily impeded both by external pressure on the lymphatic walls or by outflow resistance.

Positive pressure ventilation will push fluid from alveolus towards the interstitium and potentially towards and into the lymphatics. The positive pressure on the interstitium may compress some peripheral lymphatics potentially aiding flow, but it may also compress these thin-walled vessels impeding flow.^{93 100} A second effect is that high CVPs associated with ventilation will form a significant hydrostatic barrier to flow, given that the lymphatic pressure is usually in single figures.^{15 70 71 125} Lymph flow is again impeded.¹⁴⁷

During expiration, lower intrathoracic pressure and decreasing CVPs may allow resumption of flow, but the use of PEEP, especially high PEEP, may obviate this recovery¹¹² (Fig. 4). Positive pressure ventilation




Fig 4 Airway pressure and the lymphatics. Peak flow at a pressure of 4 mm Hg. Airway pressure of 15 mm Hg or higher compresses thin-walled collapsible lymphatics. The lymphatics drain into the central veins characterized by the superior vena cava (SVC). If the SVC pressure is high, 12 mm Hg in this example, drainage will be impeded. Hence both airway pressures and venous pressures will potentially impede drainage.

increases lung water as does PEEP.^{3 12 44 45 55 57 74 86 135} PEEP helps remove fluid from alveoli, but the reduction in thoracic duct drainage results in fluid retention in the interstitium.^{45 55 70 82 86}

In the deliberately injured lung, PEEP increases lymph production but impairs lymph flow.^{55 93 111 135} Measuring flow in the thoracic duct using a fistula shows that both positive pressure ventilation and PEEP then increase flow but, as the fistula bypasses the hydrostatic barrier of the central veins, this might be expected. Anaesthetized, ventilated animals on PEEP with high atrial pressures had less pulmonary oedema and reduced formation of pleural effusion if they had thoracic duct drainage.^{3 55} A similar mechanism was inferred in ventilated patients with pancreatitis where thoracic duct drainage was associated with an improvement in lung function.³⁷ The information available is limited, but at present there is no clear explanation of how a very low pressure system can drain effectively if there is high CVP.¹⁴⁸ The net effect of impaired drainage over time would be fluid accumulation in the lung and pleural spaces and potentially increased susceptibility to lung infection.²⁸

In the lung, there is no direct evidence linking impaired lymphatic drainage to infection risk but elsewhere, such as with chronic lymphoedema there is both fluid sequestration and predisposition to infection.^{28 79 108 117 147}

Circumstantial associations include the observation that in the critically ill nosocomial lung infection usually starts some days after commencing ventilation even in otherwise fit individuals and that time interval is commensurate with the effects of lymphatic obstruction. After lung transplant, the infection rate in the first few days is high and antibody responses are minimal.¹⁴² In these patients, lymphatic drainage re-establishes itself as early as the second week.¹⁰⁷ There is also an adverse relationship between a CVP >7 mm Hg and outcome in ventilated lung transplant patients.¹⁰¹

Ventilation impairs lymphatic drainage and chronic lymphatic obstruction predisposes to infection in other situations, but there is limited evidence for these statements.^{2 9 28}

Organ systems and positive pressure ventilation

The obvious immediate effects of positive pressure ventilation on other organs relate to a decrease in cardiac output (Fig. 5). The predominant effects on the kidney are a decrease in glomerular filtration rate and diversion of intra-renal blood flow both due to reduced cardiac output and both readily corrected either by fluid administration or by changing to a spontaneous mode of ventilation.^{5 47 103 139}

Some of the effects of PEEP on the hepato-splanchnic circulation can be reversed by fluid administration, implying a primary role for cardiac output. However, this does not apply to reduced portal blood flow or increased hepatic blood volume both of which are secondary to venous congestion, nor the reduced hepatosplanchnic lymphatic drainage.^{6 14 39 46 80 104 114 141 146} Hepatic metabolic function is impaired by PEEP of 10 mm Hg in sepsis but is less significant after cardiac surgery.^{61 114} PEEP has been shown to increase transcapillary fluid flux peripherally, thereby increasing interstitial fluid, and it also influences microcirculatory flow.^{10 57 77 96}

The sustained increases in CVP secondary to positive pressure ventilation lead to venous congestion, increased end-capillary pressure and altered fluid dynamics within organs with potentially deleterious effects.^{14 94 106 118 119} It also affects lymphatic function. In the kidney, venous congestion compresses lymphatic collecting ducts impeding drainage and a similar mechanism is postulated for both mesenteric and peritoneal drainage.^{74 75 105} Raised intra-abdominal pressure impairs lymphatic drainage and affects organ function.⁸¹ High thoracic duct pressure increases interstitial fluid in both liver and kidneys and has measurable effects on renal function.¹¹⁰ This may provide linkage to the increasing evidence that excess fluid itself may delay recovery in areas as diverse as lower gastrointestinal surgery, free flap surgery, and ventilation of acute lung injury, although this is contentious.^{22 56 62 63 109 145}

The evidence for the effects of venous and lymphatic congestion impairing organ function is as yet rudimentary, but the physics is hard to dismiss.

Conclusions

Ventilation with PEEP is a proven, effective modality in both anaesthesia and ICU, despite the immense physiological derangement it causes. The true costs incurred cannot be assessed until these physiological derangements are recognized and evaluated. The redistribution of alveolar ventilation and the associated altered capillary




Fig 5 Flow diagram illustrating the physiological interactions of positive pressure ventilation. (A) leads to (B) leads to (C)

perfusion, the functional changes in surfactant, the transcapillary fluid shifts, the impaired lymphatic drainage, and impeded venous return all contribute to wet, harder to ventilate lungs. This necessitates more aggressive ventilation, while impeded lymphatic drainage predisposes to stasis and infection. Correction of the immediate problem of reduced cardiac output by fluid administration increases central and systemic venous pressures further. The recently recognized complications of ventilation have been attributed to barotrauma and inflammation, but the other physiological mechanisms outlined here may have been ignored. Prolonged ventilation is associated with lung injury, infection, and multi-organ system dysfunction and if these may, even in part, be attributable to the physiological derangements described, then the real cost of ventilation, or oxygenation, is and has always been higher than was realized. (See Fig. 5)

There are other reasons why ventilation should be reassessed. Genuine developments in positive pressure ventilation are constrained by physics. The simple fact that only one pressure and one time-related waveform can be delivered at the tracheal tube precludes the potential for genuine technological advance. Is ventilation as a modality at an evolutionary dead end? Are there potential alternatives? It would seem logical in severe illness to avoid physiological derangement and to rest the lung. Tank or cuirasses could use the thorax to generate a negative inspiratory pressure, and maintain near-normal gradients that are fundamental to alveolar, capillary, and lymphatic function while minimizing haemodynamic disturbance. In severe illness, oxygen supplementation would inevitably be required to meet tissue requirements. Fundamental to this would be a new approach to defining tissue oxygen requirements divorced from the current approach based on measurement of arterial content. Does the technology to supplement oxygen exist? The same technological principles that revolutionized critical care dialysis from 'high tech' to bedside haemofiltration should allow the development of bedside extracorporeal oxygenation using the heart as the pump. Lung assist devices already exist, but are in their infancy.^{8 31 33 150 151}

Will maintenance of 'normal' ventilatory patterns assist recovery? The evidence from non-invasive ventilation and the occasional study using an iron lung provides circumstantial encouragement.²⁵⁻²⁷

There are many unanswered questions, but for future progress there are three preliminary steps to be taken:

- (i) recognition of the full range of physiological derangement and its potential consequences;
- (ii) accepting the physical and physiological constraints in the further evolution of positive pressure ventilation;
- (iii) acknowledgment that while positive pressure ventilation, a technology now more than 50 yr old, serves an immensely useful function, it may be time to consider alternatives in the critically ill.

What is the real cost of ventilation? We will not know until we evaluate the role of physiological derangement on late complications. It is likely the cost is high.

References

- 1 Acute Respiratory Distress Syndrome Network. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med* 2000; **342**: 1301–8
- 2 Albelda SM, Hansen-Flaschen JH, Lanken PN, Fishman AP. Effects of increased ventilation on lung lymph flow in unanesthetized sheep. *J Appl Physiol* 1986; **60**: 2063–70
- 3 Allen SJ, Drake RE, Laine GA, Gabel JC. Effect of thoracic duct drainage on hydrostatic pulmonary edema and pleural effusion in sheep. *J Appl Physiol* 1991; **71**: 314–6
- 4 Andrivet P, Adnot S, Brun-Buisson C, et al. Involvement of ANF in the acute antidiuresis during PEEP ventilation. *J Appl Physiol* 1988; **65**: 1967–74
- 5 Annat G, Viale JP, Bui Xuan B, et al. Effect of PEEP ventilation on renal function, plasma renin, aldosterone, neurophysins and urinary ADH, and prostaglandins. *Anesthesiology* 1983; **58**: 136–41
- 6 Arvidsson D, Almquist P, Haglund U. Effects of positive end-expiratory pressure on splanchnic circulation and function in experimental peritonitis. *Arch Surg* 1991; **126**: 631–6
- 7 Bark H, Le Roith D, Nyska M, Glick SM. Elevations in plasma ADH levels during PEEP ventilation in the dog: mechanisms involved. *Am J Physiol* 1980; **239**: E474–81
- 8 Bein T, Weber F, Philipp A, et al. A new pumpless extracorporeal interventional lung assist in critical hypoxemia/hypercapnia. *Crit Care Med* 2006; **34**: 1372–7
- 9 Bernard GR, Pou NA, Coggeshall JW, Carroll FE, Snapper JR. Comparison of the pulmonary dysfunction caused by cardiogenic and noncardiogenic pulmonary edema. *Chest* 1995; **108**: 798–803
- 10 Bersten AD, Gnidec AA, Rutledge FS, Sibbald WJ. Hyperdynamic sepsis modifies a PEEP-mediated redistribution in organ blood flows. *Am Rev Respir Dis* 1990; **141**: 1198–208
- 11 Bersten AD, Davidson K, Nicholas TE, Doyle IR. Respiratory mechanics and surfactant in the acute respiratory distress syndrome. *Clin Exp Pharmacol Physiol* 1998; **25**: 955–63
- 12 Blomqvist H, Frostell C, Pieper R, Hedenstierna G. Measurement of dynamic lung fluid balance in the mechanically ventilated dog. Theory and results. *Acta Anaesthesiol Scand* 1990; **34**: 370–6
- 13 Boker A, Haberman CJ, Girling L, et al. Variable ventilation improves perioperative lung function in patients undergoing abdominal aortic aneurysmectomy. *Anesthesiology* 2004; **100**: 608–16
- 14 Brienza N, Revely JP, Ayuse T, Robotham JL. Effects of PEEP on liver arterial and venous blood flows. *Am J Respir Crit Care Med* 1995; **152**: 504–10
- 15 Browse NL, Lord RS, Taylor A. Pressure waves and gradients in the canine thoracic duct. *J Physiol* 1971; **213**: 507–24
- 16 Bugge JF. Pressure limited ventilation with permissive hypoxia and nitric oxide in the treatment of adult respiratory distress syndrome. *Eur J Anaesthesiol* 1999; **16**: 799–802
- 17 Carvalho AR, Jandre FC, Pino AV, et al. Effects of descending positive end-expiratory pressure on lung mechanics and aeration in healthy anaesthetized piglets. *Crit Care* 2006; **10**: R122
- 18 Carvalho AR, Jandre FC, Pino AV, et al. Positive end-expiratory pressure at minimal respiratory elastance corresponds to the best compromise between mechanical stress and lung aeration in oleic acid-induced lung injury. *Crit Care* 2007; **11**: R86

- 19 Choudhury S, Wilson MR, Goddard ME, O'Dea KP, Takata M. Mechanisms of early pulmonary neutrophil sequestration in ventilator-induced lung injury in mice. *Am J Physiol Lung Cell Mol Physiol* 2004; **287**: L902–10
- 20 Chu EK, Whitehead T, Slutsky AS. Effects of cyclic opening and closing at low- and high-volume ventilation on bronchoalveolar lavage cytokines. *Crit Care Med* 2004; **32**: 168–74
- 21 Clark H, Reid K. The potential of recombinant surfactant protein D therapy to reduce inflammation in neonatal chronic lung disease, cystic fibrosis, and emphysema. *Arch Dis Child* 2003; **88**: 981–4
- 22 Clark JR, McCluskey SA, Hall F, et al. Predictors of morbidity following free flap reconstruction for cancer of the head and neck. *Head Neck* 2007; **29**: 1090–101
- 23 Cope DK, Parker JC, Taylor MD, Houston M, Taylor AE. Pulmonary capillary pressures during hypoxia and hypoxemia: experimental and clinical studies. *Crit Care Med* 1989; **17**: 853–7
- 24 Copland IB, Reynaud D, Pace-Asciak C, Post M. Mechanotransduction of stretch-induced prostanoid release by fetal lung epithelial cells. *Am J Physiol Lung Cell Mol Physiol* 2006; **291**: L487–95
- 25 Corrado A, Gorini M, De Paola E, et al. Iron lung treatment of acute on chronic respiratory failure: 16 yrs of experience. *Monaldi Arch Chest Dis* 1994; **49**: 552–5
- 26 Corrado A, De Paola E, Gorini M, et al. Intermittent negative pressure ventilation in the treatment of hypoxic hypercapnic coma in chronic respiratory insufficiency. *Thorax* 1996; **51**: 1077–82
- 27 Corrado A, Ginanni R, Villella G, et al. Iron lung versus conventional mechanical ventilation in acute exacerbation of COPD. *Eur Respir J* 2004; **23**: 419–24
- 28 Cui Y, Urschel JD, Petrelli NJ. The effect of cardiopulmonary lymphatic obstruction on heart and lung function. *Thorac Cardiovasc Surg* 2001; **49**: 35–40
- 29 Cunha-Goncalves D, Perez-de-Sa V, Ingimarsson J, Werner O, Larsson A. Inflation lung mechanics deteriorates markedly after saline instillation and open endotracheal suctioning in mechanically ventilated healthy piglets. *Pediatr Pulmonol* 2007; **42**: 10–4
- 30 D'Angelo E, Pecchiari M, Gentile G. Dependence of lung injury on surface tension during low-volume ventilation in normal open-chest rabbits. *J Appl Physiol* 2007; **102**: 174–82
- 31 David M, Heinrichs W. High-frequency oscillatory ventilation and an interventional lung assist device to treat hypoxaemia and hypercapnia. *Br J Anaesth* 2004; **93**: 582–6
- 32 Dehnert C, Risse F, Ley S, et al. Magnetic resonance imaging of uneven pulmonary perfusion in hypoxia in humans. *Am J Respir Crit Care Med* 2006; **174**: 1132–8
- 33 Dembinski R, Hochhausen N, Terbeck S, et al. Pumpless extracorporeal lung assist for protective mechanical ventilation in experimental lung injury. *Crit Care Med* 2007; **35**: 2359–66
- 34 Dirocco JD, Carney DE, Nieman GF. Correlation between alveolar recruitment/derecruitment and inflection points on the pressure–volume curve. *Intensive Care Med* 2007; **33**: 1204–11
- 35 dos Santos CC, Slutsky AS. Invited review: mechanisms of ventilator-induced lung injury: a perspective. *J Appl Physiol* 2000; **89**: 1645–55
- 36 dos Santos CC, Slutsky AS. The contribution of biophysical lung injury to the development of biotrauma. *Annu Rev Physiol* 2006; **68**: 585–618
- 37 Dugernier T, Reynaert MS, Deby-Dupont G, et al. Prospective evaluation of thoracic-duct drainage in the treatment of respiratory failure complicating severe acute pancreatitis. *Intensive Care Med* 1989; **15**: 372–8
- 38 Edibam C. Ventilator-induced lung injury and implications for clinical management. *Crit Care Resusc* 2000; **2**: 269–77
- 39 Elk JR, Laine GA. Pressure within the thoracic duct modulates lymph composition. *Microvasc Res* 1990; **39**: 315–21
- 40 Escamilla R, Prevost MC, Cariven C, Hermant C, Krempf M, Chap H. Bronchoalveolar lavage phospholipid abnormalities in HIV-infected patients. *Eur Respir J* 1993; **6**: 1301–7
- 41 Evgenov OV, Hevroy O, Bremnes KE, Bjertnaes LJ. Effect of aminoguanidine on lung fluid filtration after endotoxin in awake sheep. *Am J Respir Crit Care Med* 2000; **162**: 465–70
- 42 Farge D, De la Coussaye JE, Beloucif S, Fratacci MD, Payen DM. Interactions between hemodynamic and hormonal modifications during PEEP-induced antidiuresis and antinatriuresis. *Chest* 1995; **107**: 1095–100
- 43 Farias LL, Faffe DS, Xisto DG, et al. Positive end-expiratory pressure prevents lung mechanical stress caused by recruitment/derecruitment. *J Appl Physiol* 2005; **98**: 53–61
- 44 Frostell C, Blomqvist H, Pieper R, Hedenstierna G. Separate thoracic and abdominal lymph flow in the dog. *Acta Chir Scand Suppl* 1986; **530**: 5–7
- 45 Frostell C, Blomqvist H, Hedenstierna G, Halbig I, Pieper R. Thoracic and abdominal lymph drainage in relation to mechanical ventilation and PEEP. *Acta Anaesthesiol Scand* 1987; **31**: 405–12
- 46 Fujita Y. Effects of PEEP on splanchnic hemodynamics and blood volume. *Acta Anaesthesiol Scand* 1993; **37**: 427–31
- 47 Gattinoni L, Agostoni A, Damia G, et al. Hemodynamics and renal function during low frequency positive pressure ventilation with extracorporeal CO₂ removal. A comparison with continuous positive pressure ventilation. *Intensive Care Med* 1980; **6**: 155–61
- 48 Gattinoni L, Carlesso E, Cadringer P, Valenza F, Vagginelli F, Chiumello D. Physical and biological triggers of ventilator-induced lung injury and its prevention. *Eur Respir J Suppl* 2003; **47**: 15s–25s
- 49 Gattinoni L, Caironi P, Cressoni M, et al. Lung recruitment in patients with the acute respiratory distress syndrome. *N Engl J Med* 2006; **354**: 1775–86
- 50 Gehr P, Im Hof V, Geiser M, Schurch S. The mucociliary system of the lung—role of surfactants. *Schweiz Med Wochenschr* 2000; **130**: 691–8
- 51 Guyton AC, Taylor AE, Brace RA. A synthesis of interstitial fluid regulation and lymph formation. *Fed Proc* 1976; **35**: 1881–5
- 52 Guyton AC, Taylor AE, Drake RE, Parker JC. Dynamics of subatmospheric pressure in the pulmonary interstitial fluid. *Ciba Found Symp* 1976; **38**: 77–100
- 53 Haagsman HP. Interactions of surfactant protein A with pathogens. *Biochim Biophys Acta* 1998; **1408**: 264–77
- 54 Haagsman HP, Diemel RV. Surfactant-associated proteins: functions and structural variation. *Comp Biochem Physiol A Mol Integr Physiol* 2001; **129**: 91–108
- 55 Haider M, Schad H, Mendler N. Thoracic duct lymph and PEEP studies in anesthetized dogs. II. Effect of a thoracic duct fistula on the development of a hyponcotic-hydrostatic pulmonary oedema. *Intensive Care Med* 1987; **13**: 278–83
- 56 Haughey BH, Wilson E, Kluwe L, et al. Free flap reconstruction of the head and neck: analysis of 241 cases. *Otolaryngol Head Neck Surg* 2001; **125**: 10–7
- 57 Hersch M, Bersten A, Robertson S, Rutledge F, Sibbald WJ. PEEP increases non-pulmonary microvascular fluid flux in healthy and septic sheep. *Chest* 1989; **96**: 1142–9
- 58 Hinz J, Gehoff A, Moerer O, et al. Regional filling characteristics of the lungs in mechanically ventilated patients with acute lung injury. *Eur J Anaesthesiol* 2007; **24**: 414–24

- 59 Hinz J, Moerer O, Neumann P, et al. Regional pulmonary pressure volume curves in mechanically ventilated patients with acute respiratory failure measured by electrical impedance tomography. *Acta Anaesthesiol Scand* 2006; **50**: 331–9
- 60 Hinz J, Moerer O, Neumann P, Dudykevych T, Hellige G, Quintel M. Effect of positive end-expiratory-pressure on regional ventilation in patients with acute lung injury evaluated by electrical impedance tomography. *Eur J Anaesthesiol* 2005; **22**: 817–25
- 61 Holland A, Thuemer O, Schelenz C, van Hout N, Sakka SG. Positive end-expiratory pressure does not affect indocyanine green plasma disappearance rate or gastric mucosal perfusion after cardiac surgery. *Eur J Anaesthesiol* 2007; **24**: 141–7
- 62 Holte K, Klarskov B, Christensen DS, et al. Liberal versus restrictive fluid administration to improve recovery after laparoscopic cholecystectomy: a randomized, double-blind study. *Ann Surg* 2004; **240**: 892–9
- 63 Holte K, Foss NB, Andersen J, et al. Liberal or restrictive fluid administration in fast-track colonic surgery: a randomized, double-blind study. *Br J Anaesth* 2007; **99**: 500–8
- 64 Hopkins SR, Levin DL. Heterogeneous pulmonary blood flow in response to hypoxia: a risk factor for high altitude pulmonary edema? *Respir Physiol Neurobiol* 2006; **151**: 217–28
- 65 Kallet RH, Katz JA. Respiratory system mechanics in acute respiratory distress syndrome. *Respir Care Clin N Am* 2003; **9**: 297–319
- 66 Kharasch ED, Yeo KT, Kenny MA, Buffington CW. Atrial natriuretic factor may mediate the renal effects of PEEP ventilation. *Anesthesiology* 1988; **69**: 862–9
- 67 Krause MF, Wiemann T, Reisner A, Orłowska-Volk M, Kohler H, Ankermann T. Surfactant reduces extravascular lung water and invasion of polymorphonuclear leukocytes into the lung in a piglet model of airway lavage. *Pulm Pharmacol Ther* 2005; **18**: 129–39
- 68 Kredel M, Muellenbach RM, Brock RW, et al. Liver dysfunction after lung recruitment manoeuvres during pressure-controlled ventilation in experimental acute respiratory distress. *Crit Care* 2007; **11**: R13
- 69 Lachmann RA, van Kaam AH, Haitsma JJ, Lachmann B. High positive end-expiratory pressure levels promote bacterial translocation in experimental pneumonia. *Intensive Care Med* 2007; **33**: 1800–4
- 70 Laine GA, Allen SJ, Katz J, Gabel JC, Drake RE. Effect of systemic venous pressure elevation on lymph flow and lung edema formation. *J Appl Physiol* 1986; **61**: 1634–8
- 71 Laine GA, Allen SJ, Katz J, Gabel JC, Drake RE. Outflow pressure reduces lymph flow rate from various tissues. *Microvasc Res* 1987; **33**: 135–42
- 72 Lassen HC. A preliminary report on the 1952 epidemic of poliomyelitis in Copenhagen with special reference to the treatment of acute respiratory insufficiency. *Lancet* 1953; **1**: 37–41
- 73 Lassen HC, Bjerneboe M, Ibsen B, Neukirch F. Treatment of tetanus with curarisation, general anaesthesia, and intratracheal positive-pressure ventilation. *Lancet* 1954; **267**: 1040–4
- 74 Last M, Kurtz L, Stein TA, Wise L. Effect of PEEP on the rate of thoracic duct lymph flow and clearance of bacteria from the peritoneal cavity. *Am J Surg* 1983; **145**: 126–30
- 75 Lattuada M, Hedenstierna G. Abdominal lymph flow in an endotoxin sepsis model: influence of spontaneous breathing and mechanical ventilation. *Crit Care Med* 2006; **34**: 2792–8
- 76 Lebel L, Smith L, Risberg B, Gerdin B, Laurent TC. Effect of increased hydrostatic pressure on lymphatic elimination of hyaluronan from sheep lung. *J Appl Physiol* 1988; **64**: 1327–32
- 77 Lehtipalo S, Biber B, Frojse R, Arnerlov C, Johansson G, Winso O. Effects of positive end-expiratory pressure on intestinal circulation during graded mesenteric artery occlusion. *Acta Anaesthesiol Scand* 2001; **45**: 875–84
- 78 Lin CY, Zhang H, Cheng KC, Slutsky AS. Mechanical ventilation may increase susceptibility to the development of bacteremia. *Crit Care Med* 2003; **31**: 1429–34
- 79 Lopez FA, Lartchenko S. Skin and soft tissue infections. *Infect Dis Clin North Am* 2006; **20**: 759–72
- 80 Love R, Choe E, Lippton H, Flint L, Steinberg S. Positive end-expiratory pressure decreases mesenteric blood flow despite normalization of cardiac output. *J Trauma* 1995; **39**: 195–9
- 81 Malbrain ML, Pelosi P, De laet I, Lattuada M, Hedenstierna G. Lymphatic drainage between thorax and abdomen: please take good care of this well-performing machinery. *Acta Clin Belg Suppl* 2007; **11**: 152–61
- 82 Mallick A, Bodenham AR. Disorders of the lymph circulation: their relevance to anaesthesia and intensive care. *Br J Anaesth* 2003; **91**: 265–72
- 83 Malloy JL, Veldhuizen RA, Lewis JF. Effects of ventilation on the surfactant system in sepsis-induced lung injury. *J Appl Physiol* 2000; **88**: 401–8
- 84 Mandava S, Kolobow T, Vitale G, et al. Lethal systemic capillary leak syndrome associated with severe ventilator-induced lung injury: an experimental study. *Crit Care Med* 2003; **31**: 885–92
- 85 Marini JJ. Relative importance of stretch and shear in ventilator-induced lung injury. *Crit Care Med* 2004; **32**: 302–4
- 86 Maybauer DM, Talke PO, Westphal M, et al. Positive end-expiratory pressure ventilation increases extravascular lung water due to a decrease in lung lymph flow. *Anaesth Intensive Care* 2006; **34**: 329–33
- 87 Meier T, Luepschen H, Karsten J, et al. Assessment of regional lung recruitment and derecruitment during a PEEP trial based on electrical impedance tomography. *Intensive Care Med* 2008; **34**: 543–50
- 88 Mergoni M, Rossi A. Physiopathology of acute respiratory failure in COPD and asthma. *Minerva Anestesiol* 2001; **67**: 198–205
- 89 Milic-Emili J, Henderson JA, Dolovich MB, Trop D, Kaneko K. Regional distribution of inspired gas in the lung. *J Appl Physiol* 1966; **21**: 749–59
- 90 Miranda DR, Klompe L, Mekel J, et al. Open lung ventilation does not increase right ventricular outflow impedance: an echo-Doppler study. *Crit Care Med* 2006; **34**: 2555–60
- 91 Mitaka C, Nagura T, Sakanishi N, Tsunoda Y, Amaha K. Two-dimensional echocardiographic evaluation of inferior vena cava, right ventricle, and left ventricle during positive-pressure ventilation with varying levels of positive end-expiratory pressure. *Crit Care Med* 1989; **17**: 205–10
- 92 Moloney ED, Griffiths MJ. Protective ventilation of patients with acute respiratory distress syndrome. *Br J Anaesth* 2004; **92**: 261–70
- 93 Moriondo A, Mukenge S, Negrini D. Transmural pressure in rat initial subpleural lymphatics during spontaneous or mechanical ventilation. *Am J Physiol Heart Circ Physiol* 2005; **289**: H263–9
- 94 Mortillaro NA, Taylor AE. Interstitial fluid pressure of ileum measured from chronically implanted polyethylene capsules. *Am J Physiol* 1989; **257**: H62–9
- 95 Mutch WA, Eschun GM, Kowalski SE, Graham MR, Girling LG, Lefevre GR. Biologically variable ventilation prevents deterioration of gas exchange during prolonged anaesthesia. *Br J Anaesth* 2000; **84**: 197–203
- 96 Mutlu GM, Mutlu EA, Factor P. GI complications in patients receiving mechanical ventilation. *Chest* 2001; **119**: 1222–41
- 97 Nieman GF, Paskanik AM, Bredenberg CE. Effect of positive end-expiratory pressure on alveolar capillary perfusion. *J Thorac Cardiovasc Surg* 1988; **95**: 712–6

- 98 Niermeyer S, Yang P, Shanmina , Drolkar , Zhuang J, Moore LG. Arterial oxygen saturation in Tibetan and Han infants born in Lhasa, Tibet. *N Engl J Med* 1995; **333**: 1248–52
- 99 Panda AK, Nag K, Harbottle RR, et al. Effect of acute lung injury on structure and function of pulmonary surfactant films. *Am J Respir Cell Mol Biol* 2004; **30**: 641–50
- 100 Pearse DB, Searcy RM, Mitzner W, Permutt S, Sylvester JT. Effects of tidal volume and respiratory frequency on lung lymph flow. *J Appl Physiol* 2005; **99**: 556–63
- 101 Pilcher DV, Scheinkestel CD, Snell GI, Davey-Quinn A, Bailey MJ, Williams TJ. High central venous pressure is associated with prolonged mechanical ventilation and increased mortality after lung transplantation. *J Thorac Cardiovasc Surg* 2005; **129**: 912–8
- 102 Pinsky MR. Cardiovascular issues in respiratory care. *Chest* 2005; **128**: 592S–7S
- 103 Priebe HJ, Heimann JC, Hedley-Whyte J. Mechanisms of renal dysfunction during positive end-expiratory pressure ventilation. *J Appl Physiol* 1981; **50**: 643–9
- 104 Risoe C, Hall C, Smiseth OA. Splanchnic vascular capacitance and positive end-expiratory pressure in dogs. *J Appl Physiol* 1991; **70**: 818–24
- 105 Rohn DA, Stewart RH, Elk JR, Laine GA, Drake RE. Renal lymphatic function following venous pressure elevation. *Lymphology* 1996; **29**: 67–75
- 106 Rossaint R, Krebs M, Forther J, Unger V, Falke K, Kaczmarczyk G. Inferior vena caval pressure increase contributes to sodium and water retention during PEEP in awake dogs. *J Appl Physiol* 1993; **75**: 2484–92
- 107 Ruggiero R, Muz J, Fietsam R, Jr, et al. Reestablishment of lymphatic drainage after canine lung transplantation. *J Thorac Cardiovasc Surg* 1993; **106**: 167–71
- 108 Ruocco V, Schwartz RA, Ruocco E. Lymphedema: an immunologically vulnerable site for development of neoplasms. *J Am Acad Dermatol* 2002; **47**: 124–7
- 109 Sakr Y, Vincent JL, Reinhart K, et al. High tidal volume and positive fluid balance are associated with worse outcome in acute lung injury. *Chest* 2005; **128**: 3098–108
- 110 Satta A, Contu B, Branca GF, Menghi R, Bresadola F, Bartoli E. Importance of liver interstitial pressure on sodium retention. *Nephron* 1988; **49**: 190–6
- 111 Schad H, Flowaczny H, Brechtelsbauer H, Birkenfeld G. The significance of respiration for thoracic duct flow in relation to other driving forces of lymph flow. *Pflugers Arch* 1978; **378**: 121–5
- 112 Schmid-Schonbein GW. Microlymphatics and lymph flow. *Physiol Rev* 1990; **70**: 987–1028
- 113 Schmidt R, Markart P, Ruppert C, et al. Pulmonary surfactant in patients with Pneumocystis pneumonia and acquired immunodeficiency syndrome. *Crit Care Med* 2006; **34**: 2370–6
- 114 Schrickler T, Kugler B, Schywalsky M, Braun G, Trager K, Georgieff M. Effects of PEEP ventilation on liver metabolism. *Infusionsther Transfusionsmed* 1995; **22**: 168–74
- 115 Seeger W, Pison U, Buchhorn R, Obertacke U, Joka T. Surfactant abnormalities and adult respiratory failure. *Lung* 1990; **168**: 891–902
- 116 Seeger W, Gunther A, Walrath HD, Grimminger F, Lasch HG. Alveolar surfactant and adult respiratory distress syndrome. Pathogenetic role and therapeutic prospects. *Clin Invest* 1993; **71**: 177–90
- 117 Serpell JW, Carne PW, Bailey M. Radical lymph node dissection for melanoma. *ANZ J Surg* 2003; **73**: 294–9
- 118 Shibayama Y. The role of hepatic venous congestion and endotoxaemia in the production of fulminant hepatic failure secondary to congestive heart failure. *J Pathol* 1987; **151**: 133–8
- 119 Shinozaki M. Respiratory and cardiovascular management of septic ALI-ARDS and shock. *Nippon Rinsho* 2004; **62**: 2301–7
- 120 Siegel LC, Pearl RG, August DA. Pulmonary capillary pressure measurement during global hypoxia in sheep. *Anesth Analg* 1993; **76**: 149–55
- 121 Simbruner G, Mittal RA, Smith J, et al. Effects of duration and amount of lung stretch at biophysical, biochemical, histological, and transcriptional levels in an in vivo rabbit model of mild lung injury. *Am J Perinatol* 2007; **24**: 149–59
- 122 Slutsky AS, Imai Y. Ventilator-induced lung injury, cytokines, PEEP, and mortality: implications for practice and for clinical trials. *Intensive Care Med* 2003; **29**: 1218–21
- 123 Steinberg JM, Schiller HJ, Halter JM, et al. Alveolar instability causes early ventilator-induced lung injury independent of neutrophils. *Am J Respir Crit Care Med* 2004; **169**: 57–63
- 124 Suter PM. Reducing ventilator-induced lung injury and other organ injury by the prone position. *Crit Care* 2006; **10**: 139
- 125 Szabo G, Magyar Z. Effect of increased systemic venous pressure on lymph pressure and flow. *Am J Physiol* 1967; **212**: 1469–74
- 126 Taskar V, John J, Evander E, Robertson B, Jonson B. Surfactant dysfunction makes lungs vulnerable to repetitive collapse and reexpansion. *Am J Respir Crit Care Med* 1997; **155**: 313–20
- 127 Terragni PP, Rosboch G, Tealdi A, et al. Tidal hyperinflation during low tidal volume ventilation in acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2007; **175**: 160–6
- 128 Tremblay LN, Miatto D, Hamid Q, Govindarajan A, Slutsky AS. Injurious ventilation induces widespread pulmonary epithelial expression of tumor necrosis factor-alpha and interleukin-6 messenger RNA. *Crit Care Med* 2002; **30**: 1693–700
- 129 Turunen R, Nupponen I, Siitonen S, Repo H, Andersson S. Onset of mechanical ventilation is associated with rapid activation of circulating phagocytes in preterm infants. *Pediatrics* 2006; **117**: 448–54
- 130 Tusman G, Suarez-Sipmann F, Bohm SH, et al. Monitoring dead space during recruitment and PEEP titration in an experimental model. *Intensive Care Med* 2006; **32**: 1863–71
- 131 Ueda T, Ikegami M. Change in properties of exogenous surfactant in injured rabbit lung. *Am J Respir Crit Care Med* 1996; **153**: 1844–9
- 132 Uhlig S. Ventilation-induced lung injury and mechanotransduction: stretching it too far? *Am J Physiol Lung Cell Mol Physiol* 2002; **282**: L892–6
- 133 Valente Barbas CS. Lung recruitment maneuvers in acute respiratory distress syndrome and facilitating resolution. *Crit Care Med* 2003; **31**: S265–71
- 134 van den Berg PC, Grimbergen CA, Spaan JA, Pinsky MR. Positive pressure inspiration differentially affects right and left ventricular outputs in postoperative cardiac surgery patients. *J Crit Care* 1997; **12**: 56–65
- 135 van der Zee H, Cooper JA, Hakim TS, Malik AB. Alterations in pulmonary fluid balance induced by positive end-expiratory pressure. *Respir Physiol* 1986; **64**: 125–33
- 136 Vanderzwan J, McCaig L, Mehta S, et al. Characterizing alterations in the pulmonary surfactant system in a rat model of *Pseudomonas aeruginosa* pneumonia. *Eur Respir J* 1998; **12**: 1388–96
- 137 Veldhuizen RA, Tremblay LN, Govindarajan A, van Roozendaal BA, Haagsman HP, Slutsky AS. Pulmonary surfactant is altered during mechanical ventilation of isolated rat lung. *Crit Care Med* 2000; **28**: 2545–51
- 138 Veldhuizen RA, Slutsky AS, Joseph M, McCaig L. Effects of mechanical ventilation of isolated mouse lungs on surfactant and inflammatory cytokines. *Eur Respir J* 2001; **17**: 488–94
- 139 Venus B, Mathru M, Smith RA, Pham CG, Shirakawa Y, Sugiura A. Renal function during application of positive end-expiratory

- pressure in swine: effects of hydration. *Anesthesiology* 1985; **62**: 765–9
- 140** Wagner WW Jr. Capillary recruitment in the pulmonary microcirculation. *Chest* 1988; **93**: 85S–8S
- 141** Walfisch S, Weksler N, Fisher A, Shapira Y. Effects of oleic acid lung injury and positive end-expiratory pressure on central hemodynamics and regional blood flow. *Isr J Med Sci* 1997; **33**: 14–7
- 142** Wang F, Winter JB, Dam M, Wildevuur CR, Prop J. Influence of interrupted pulmonary lymph drainage on antibody responses in hilar-stripped lungs. *J Heart Lung Transplant* 1992; **11**: S215–20
- 143** West JB. Invited review: pulmonary capillary stress failure. *J Appl Physiol* 2000; **89**: 2483–9
- 144** West JB. The physiological challenges of the 1952 Copenhagen poliomyelitis epidemic and a renaissance in clinical respiratory physiology. *J Appl Physiol* 2005; **99**: 424–32
- 145** Wiedemann HP, Wheeler AP, Bernard GR, et al. Comparison of two fluid-management strategies in acute lung injury. *N Engl J Med* 2006; **354**: 2564–75
- 146** Winso O, Biber B, Gustavsson B, Holm C, Milsom I, Niemand D. Portal blood flow in man during graded positive end-expiratory pressure ventilation. *Intensive Care Med* 1986; **12**: 80–5
- 147** Witte CL, Witte MH. Disorders of lymph flow. *Acad Radiol* 1995; **2**: 324–34
- 148** Wu J, Mack GW. Effect of lymphatic outflow pressure on lymphatic albumin transport in humans. *J Appl Physiol* 2001; **91**: 1223–8
- 149** Zhang H, Downey GP, Suter PM, Slutsky AS, Ranieri VM. Conventional mechanical ventilation is associated with bronchoalveolar lavage-induced activation of polymorphonuclear leukocytes: a possible mechanism to explain the systemic consequences of ventilator-induced lung injury in patients with ARDS. *Anesthesiology* 2002; **97**: 1426–33
- 150** Zimmermann M, Bein T, Philipp A, et al. Interhospital transportation of patients with severe lung failure on pumpless extracorporeal lung assist. *Br J Anaesth* 2006; **96**: 63–6
- 151** Zimmermann M, Philipp A, Schmid FX, Dorlac W, Arlt M, Bein T. From Baghdad to Germany: use of a new pumpless extracorporeal lung assist system in two severely injured US soldiers. *Asaio J* 2007; **53**: e4–6